

Evaluation of Obstetric Outcomes in Women with Endometriosis

Endometriozisli Kadınlarda Obstetrik Sonuçların Değerlendirilmesi

^{1b}Ufuk ATLIHAN^a, ^{1b}Feruh ACET^b, ^{1b}Onur YAVUZ^c, ^{1b}Burak ERSAK^d, ^{1b}Can ATA^e,
^{1b}Tevfik Berk BİLDACI^f, ^{1b}Selçuk ERKILINÇ^f, ^{1b}Hüseyin Aytuğ AVŞAR^g

^aPrivate Karataş Hospital Gynecology and Obstetrics Clinic, İzmir, Türkiye

^bEge University Faculty of Medicine, Health Application and Research Center, Department of Gynecology and Obstetrics, İzmir, Türkiye

^cDokuz Eylül University Faculty of Medicine, Department of Gynecology and Obstetrics, İzmir, Türkiye

^dAnkara Etilik City Hospital Gynecology and Obstetrics Clinic, Ankara, Türkiye

^eBuca Seyfi Demirsöy Training and Research Hospital Gynecology and Obstetrics Clinic, İzmir, Türkiye

^fİzmir Democracy University Faculty of Medicine, Buca Seyfi Demirsöy Training and Research Hospital, Department of Gynecology and Obstetrics, İzmir, Türkiye

^gİzmir Tinaztepe University Private Galen Hospital, Department of Gynecology and Obstetrics, İzmir, Türkiye

ABSTRACT

Objective: It is generally accepted that the endometrium of women with endometriosis is abnormal, although there is ongoing debate as to whether these abnormalities impair decidualization and placentation during pregnancy. The aim of this study is to evaluate the obstetric and neonatal outcomes in patients diagnosed with endometriosis. **Material and Methods:** 1015 patients who underwent pregnancy follow-up in our obstetrics clinic and gave birth in our hospital between 2018 and 2023 were retrospectively examined. The patients evaluated in the study were evaluated in two separate groups according to the presence of endometriosis. The presence of preterm delivery, gestational diabetes (GDM), gestational hypertension (GHT), preeclampsia, premature rupture of membranes (PROM), fetal growth restriction (FGR), Neonatal Intensive Care Unit (NICU) were evaluated in all patients. **Results:** ART presence was found to be significantly higher in the endometriosis group ($p=0.038$). Gestational week was found to be significantly lower in the endometriosis group ($p=0.018$). The GHT presence was found to be significantly higher in the endometriosis group ($p=0.034$). The Cesarean presence was found to be significantly higher in the endometriosis group ($p=0.037$). Estimated blood loss volume was significantly higher in the endometriosis group ($p=0.042$). The NICU rate was significantly higher in the endometriosis group ($p=0.044$). **Conclusion:** Perinatal and neonatal outcomes resulting from endometriosis depend on multifactorial factors. Prospective and large population-based studies or meta-analyses are needed to clarify possible risks.

Keywords: Endometriosis; neonatal; perinatal

ÖZET

Amaç: Endometriozisli kadınların endometriumunun anormal olduğu genel olarak kabul edilmektedir, ancak bu anormalliklerin gebelik sırasında desidualizasyon ve plasentasyona zarar verip vermediği konusunda tartışmalar devam etmektedir. Bu çalışmanın amacı endometriozis tanısı almış hastalarda obstetrik ve neonatal sonuçları değerlendirmektir. **Gereç ve Yöntemler:** 2018-2023 yılları arasında kadın doğum kliniğimizde gebelik takibi yapılan ve hastanemizde doğum yapan 1015 hasta retrospektif olarak incelendi. Çalışmada değerlendirilen hastalar endometriozis varlığına göre iki ayrı grupta değerlendirildi. Tüm hastalarda preterm doğum, gestasyonel diyabet (GDM), gestasyonel hipertansiyon (GHT), preeklampsi, erken membran rüptürü (PROM), fetal büyüme kısıtlaması (FGR), Yenidoğan Yoğun Bakım Ünitesi (NICU) varlığı değerlendirildi. **Bulgular:** ART öyküsünün endometriozis grubunda anlamlı olarak daha yüksek saptandı ($p=0.038$). Gebelik haftasının endometriozis grubunda anlamlı olarak daha düşük olduğu görüldü ($p=0.018$). GHT oranının endometriozis grubunda anlamlı olarak daha yüksek saptandı ($p=0.034$). Sezaryen oranının endometriozis grubunda anlamlı olarak daha yüksek saptandı ($p=0.037$). Tahmini kan kaybı hacmi endometriozis grubunda anlamlı olarak daha yüksek saptandı ($p=0.042$). NICU oranı endometriozis grubunda anlamlı olarak daha yüksek saptandı ($p=0.044$). **Sonuç:** Endometriozis kaynaklı perinatal ve neonatal sonuçlar multifaktöriyel faktörlere bağlıdır. Olası riskleri açıklığa kavuşturmak için prospektif ve geniş popülasyon tabanlı çalışmalara veya meta-analizlere ihtiyaç vardır.

Anahtar Kelimeler: Endometriozis; neonatal; perinatal

Correspondence: Ufuk ATLIHAN

Private Karataş Hospital Gynecology and Obstetrics Clinic, İzmir, Türkiye

E-mail: cfl.ufuk@gmail.com



Peer review under responsibility of Turkish Journal of Reproductive Medicine and Surgery.

Received: 12 Sep 2024

Accepted: 05 Dec 2024

Available online: 11 Dec 2024

2587-0084 / Copyright © 2025 by Reproductive Medicine, Surgical Education, Research and Practice Foundation.
This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

Endometriosis is a chronic inflammatory disease defined as the presence of endometrial-like tissue outside the uterine cavity, and common symptoms include dysmenorrhea, dyspareunia, and subfertility.¹⁻³ Endometriosis is considered a benign gynecological disease that affects 10% of women of reproductive age throughout their lives and has an increasing incidence.⁴ The prevalence of deep and ovarian endometriosis in pregnancy is approximately 5%, which is similar to the prevalence in women attending a general gynecology clinic, and approximately 50% of women are unaware that they have the condition.^{5,6} It is generally accepted that the endometrium of women with endometriosis is abnormal, but there is ongoing debate as to whether these abnormalities impair decidualization and placentation during pregnancy.^{7,8} Since these processes may be critical for pregnancy implantation and development, it has been hypothesized that pregnancy outcome in women with endometriosis may be affected.⁹ Epidemiological studies have shown that women with some types of endometriosis may have an increased risk of preterm delivery and small for gestational age (SGA). The reason for this is changes in endometrial functions and prostaglandin (PG) levels.^{10,11} Studies on the relationship between endometriosis and pregnancy outcome are contradictory. While literature reports an increased risk of preterm delivery, preeclampsia, and prenatal hemorrhage/placental complications, some studies have not found a relationship.¹²⁻¹⁶ Previous studies reporting on obstetric complications in women with endometriosis have been based on fertility populations, retrospective data, or national statistics; the true complication rate in women with endometriosis is unknown.¹⁷⁻²⁰ The aim of this study is to evaluate the obstetric and neonatal outcomes in patients diagnosed with endometriosis.

MATERIALS AND METHODS

In present study was designed as a retrospective multicentric cohort study. The study was designed according to the Helsinki Declaration and informed consent forms were obtained from all patients. The study was initiated after receiving ethics committee approval numbered 24/26-6 from the hospital ethics committee. 1015 patients who underwent pregnancy follow-up in our obstetrics clinic and gave birth be-

tween 2018 and 2023 were retrospectively examined. The patients evaluated in the study were evaluated in two separate groups according to the presence of endometriosis. Pelvic or transvaginal ultrasonography data of all patients were evaluated from the hospital database for the presence of congenital and acquired uterine pathologies, including adenomyosis, uterine fibroids, and congenital uterine anomalies in addition to endometriosis. The presence of adenomyosis was defined according to the morphological uterus sonographic assessment criteria.²¹ Myomas were defined as well-defined masses with posterior shadowing and circumferential vascularity within or communicating with the myometrium of the deuterine body or cervix in the first trimester of pregnancy.²² Congenital uterine anomalies were evaluated according to the current ASRM classification system.²³ Ultrasonography data of the anterior and posterior pelvic compartments were evaluated for the presence of bilateral adnexal endometrioma and deep endometriosis. The International Deep Endometriosis Analysis Group accepted the presence of endometriosis as histologically proven endometriosis by surgery or the presence of lesions on ultrasound.²⁴ The presence of type of delivery, preterm delivery, gestational diabetes (GDM), gestational hypertension (GHT), assisted reproductive technology (ART) preeclampsia, premature rupture of membranes (PROM), fetal growth restriction (FGR), Neonatal Intensive Care Unit (NICU) were evaluated in all patients. The American Diabetes Association Criteria were used to diagnose gestational diabetes.²⁵ In the diagnosis of GDM, fasting blood glucose value > 92 mg/dL, first hour blood glucose value > 180 mg/dL, second hour blood glucose value > 153 mg/dL were determined as criteria. Diagnosis was made if any of the current values were exceeded. In women with no known history of diabetes mellitus, a 75-g OGTT test is performed at 24-28 weeks to measure fasting, 1-hour and 2-hour plasma glucose. Fasting for >8 hours is recommended for optimal evaluation of OGTT results.²⁵ GHT was diagnosed in accordance with the most recent American College of Obstetricians and Gynecologists bulletin.²⁶ The combination of hypertension and proteinuria is used for the diagnosis of preeclampsia. GHT is defined as blood pressure lev-

els of at >140 mm Hg as systolic or at >90 mm Hg as diastolic in measurements taken four hours or longer > 20th week of pregnancy in a woman whose blood pressure values were previously normal. Severe hypertension is considered when blood pressure is at >160 mm Hg systolic or at >110 mm Hg diastolic. To diagnose preeclampsia, women with hypertension also require the presence of proteinuria, defined as at >300 mg in a 24-hour urine collection. GHT is diagnosed in patients who meet hypertension criteria for preeclampsia without proteinuria or serious additional problems.²⁶ The Delphi Criteria were used to diagnose FGR.²⁷ When considering the Delphi FGR criteria, two single parameters (abdominal circumference (AC) or EFW < 3%) can be taken into account. Alternatively, cumulative evaluation of 4 parameters has been suggested. In these criteria (EFW or AC < 10th percentile): AC or EFW exceeding the percentiles in growth charts by > two quartiles and cerebroplacental ratio <5% or UA-PI >95% were accepted.²⁷ The estimated volume of blood loss was measured by utilizing the pregnant women's height, weight, and prenatal and postnatal Hct values.²⁸ Blood transfusion indications were determined in terms of the vital signs, estimated blood loss volume, and postpartum Hb value <8 g/dL.²⁸

STATISTICAL ANALYSIS

Statistical analysis was conducted by utilizing the SPSS 26.0 (IBM-Inc.-Chicago-IL-USA). The normality of the distribution was evaluated with the Kol-

mogrov-Smirnov Test. Mean±Standard Deviation (SD) was used for evaluating normally distributed data, and median (range) was used for non-normally distributed data. Number (n) and percentage (%) were used for evaluating categorical data. The Fisher's Exact were employed in the categorical data analysis. Logistic regression analysis was used to determine the presence of endometriosis and adverse perinatal and neonatal outcomes.

RESULTS

ART history was found to be 7.1% in the endometriosis group and 4.4% in the group without endometriosis, and was found to be significantly higher in the endometriosis group (0.038). The mean gestational age at delivery was 38+4 weeks in the endometriosis group and 39+3 weeks in the group without endometriosis, and was found to be significantly lower in the endometriosis group (p=0.018) (Table 1).

GHT rate was found to be 6.1% in the endometriosis group and 2.4% in the non-endometriosis group, and it was found to be significantly higher in the endometriosis group (p=0.034). C/S rate was found to be 46.6% in the endometriosis group and 38.5% in the non-endometriosis group, and it was found to be significantly higher in the endometriosis group (p=0.037). The estimated blood loss volume level was 545 (210-870) cc in the endometriosis group and 340 (120-650) cc in the non-endometriosis group, and was significantly higher in the endometriosis group (p=0.042) (Table 2).

TABLE 1: Comparison of demographic and obstetric data of patients.

	Endometriosis (+) n=210	Endometriosis (-) n=805	p
	median (min-max)		
Age (year)	32 (20-42)	30 (18-44)	0.38
BMI (kg/m ²)	24.1(18.2-35.6)	24.5 (18.4-34.8)	0.44
Smoking, n (%)	20 (%9.5)	81 (%10.1)	0.28
Gravidity	1.8 (1-4)	1.9 (1-4)	0.66
Parity	1.6 (1-4)	1.7 (1-4)	0.56
Multiple pregnancy, n (%)	14(%6.6)	56(%6.9)	0.61
ART, n (%)	15 (%7.1)	36(%4.4)	0.038
Abortion, n (%)	16 (%7.6)	57 (%7)	0.34
Gestational week	38+4 (31-41)	39+3 (29-41)	0.018

BMI: Body mass index, ART: Assisted reproductive technology.

TABLE 2: Comparison of perinatal outcomes between groups

	Endometriosis (+) n=210	Endometriosis (-) n=805	OR (%95 CI)	p
	n (%)	n (%)		
GHT	13 (%6.1)	20 (%2.4)	2.54 (1.22–8.08)	0.034
Preeclampsia	9 (%4.2)	17 (%2.1)	2.02(0.62–8.34)	0.26
GDM	18 (%8.5)	66 (%8.1)	1.05(0.54–2.22)	0.88
Vaginal birth	112 (%53.4)	495 (%61.5)	1.21(1.09–2.02)	0.037
C/S	98 (%46.6)	310 (%38.5)		
Preterm birth	14 (%6.6)	34 (%4.2)	1.57(0.92–3.92)	0.076
PROM	13 (%6.1)	35 (%4.3)	1.42(0.9–3.82)	0.088
FGR	12 (%5.7)	47 (%5.8)	0.98(0.88–3.54)	0.90
ΔHb	0.8±0.6	0.6±0.7	1.33(1.12–2.54)	0.14
Estimated blood loss volume (cc)	545 (210-870)	340 (120-650)	1.60(1.37–3.26)	0.042
Blood transfusion	16 (%7.8)	44 (%5.4)	1.44(1.18–2.96)	0.1

GHT: Gestational hypertension, GDM: Gestational diabetes mellitus, C/S: Cesarean section, PROM: Premature rupture of membranes, FGR: Fetal growth restriction, ΔHb: Preop-Postop difference

TABLE 3: Comparison of neonatal outcomes between groups.

	Endometriosis (+) n=210	Endometriosis (-) n=805	OR (%95 CI)	p
	Mean±SD	Mean±SD		
Apgar (1 st min)	8.2±0.8	7.8±0.7	1.05 (0.92–1.09)	0.11
Apgar (5 th min)	8.7±1.1	8.3±0.9	1.04(0.90–1.16)	0.16
Birth weight (gr)	3070±580	3190±640	0.96(0.84–1.18)	0.66
NICU, n (%)	18 (%8.4)	31 (%3.8)	2.21(1.14–4.08)	0.044

NICU: Neonatal Intensive Care Unit

NICU rate was found to be 8.4% in the endometriosis group and 3.8% in the non-endometriosis group, and it was found to be significantly higher in the endometriosis group (p=0.044) (Table 3).

DISCUSSION

In present study showed that in the group with endometriosis, in terms of pregnancy outcomes, the GHT rate, C/S rate, NICU requirement and, estimated blood loss volume were significantly higher in the endometriosis group. Studies evaluating the perinatal and neonatal effects of endometriosis in the literature have presented results from a very broad perspective. We believe that these differences between the results are due to many factors such as patient selection, number of patients, diagnostic method of endometriosis, severity of endometriosis and study methodology. The relationship between endometriosis

and preterm delivery is one of the perinatal complications that is particularly emphasized. It has been reported that the eutopic endometrium and the junctional zone are abnormal at the molecular and functional levels, leading to impaired endometrial growth, maturation and decidualization, endometrial receptivity, defective spiral artery remodeling, and defective deep placentation.^{8,9} Defective arterial remodeling is associated with a number of pregnancy complications, including preterm delivery, preeclampsia and FGR.^{9,29} Abnormal placentation may increase the risk of placental complications during pregnancy. Endometriosis is also associated with a chronic pelvic inflammatory process, and increased levels of prostaglandins and cytokines have been documented in the peritoneal fluid of women with Endometriosis.³⁰⁻³³ Increased levels of these proinflammatory mediators may stimulate myome-

trial contractions and cervical ripening, leading to preterm delivery.^{12,34} In addition, the normal frequency and amplitude of uterine contractions are altered in women with endometriosis, which may affect embryo delivery and implantation.^{35,36} Chronic inflammation may constitute the biochemical background for preterm delivery in women with endometriosis. A meta-analysis by Lalani et al. showed that the probability of preterm delivery is higher in women diagnosed with pelvic Endometriosis.³⁷ The meta-analysis by Breintoft et al. similarly showed that the probability of preterm delivery was higher in women diagnosed with pelvic endometriosis.³⁸ A study by Exacoustos et al. found a correlation between the endometriosis and preterm delivery.³⁹ The study by Farella et al. showed a higher prevalence of preterm delivery in women with a history of surgical treatment for endometriosis, especially in those with deep disease of the rectum or bladder.⁴⁰ A study on endometriosis and adverse obstetric outcomes based on more than 1.4 million births in Sweden found that endometriosis was associated with preterm delivery.¹² However, Aris, et al. reported in their study that women with endometriosis had no increased risk of preterm delivery.⁴¹ Similarly, Mekaru, et al. found in their study that women with endometriosis had no increased risk of preterm delivery.¹⁶ Although the preterm delivery rate was found to be higher in the endometriosis group, no statistically difference was found between the groups. Theories that may support excessive blood loss during cesarean section include a number of associations with angiogenesis, mild bleeding disorders, pelvic adhesions, surgical complexity, increased operative time, or bleeding from endometriotic foci.^{42,43} Endometriotic lesions may be more prone to bleeding when disturbed during pregnancy and surgery.⁴³ Decidualization of endometriotic lesions is a hormonally induced phenomenon that women with endometriosis during pregnancy.⁵ Stromal vascularity, immune cell influx, and edema from lesions may also contribute to intraoperative blood loss.^{44,45} Some women who experience excessive intraperitoneal bleeding at ovulation are at increased risk of developing deep endometriosis, but if the bleeding disorder

is clinically significant, we would also expect excessive blood loss during vaginal delivery.⁴⁶ In the study by Lafleur et al., active endometriosis was associated with an increased risk of severe hemorrhage, whereas inactive endometriosis was less strongly associated.⁴⁷ In the meta-analysis by Breintoft et al., no significant difference was observed in the frequency of postpartum bleeding in women diagnosed with endometriosis compared to women without endometriosis.³⁸ In our current study, while no significant difference was found between the groups in terms of the presence of massive bleeding requiring blood transfusion, the estimated blood loss volume level was found to be significantly higher in the endometriosis group. In the nationwide study by Stephansson et al., no association was observed between SGA and endometriosis.¹² Similarly, no association was found between endometriosis and SGA in the Danish cohort study by Glavind et al.⁴⁸ Fernando et al. suggested that women with endometrioma have an increased risk of SGA, but this risk is not present in women with other forms of endometriosis.¹⁰ In our results, no significant difference was found in terms of FGR rate between the endometriosis group and the non-endometriosis group. The relationship between endometriosis and gestational hypertension and preeclampsia is another issue that has been emphasized in the literature, with conflicting results. In the study by Chen et al., no association was found between the presence of endometriosis and the risk of gestational hypertension.⁴⁹ In the study conducted by Farland et al., women with a history of laparoscopically confirmed endometriosis had a 30% higher risk of developing hypertensive disorders during pregnancy.⁵⁰ In the study by Hadfield et al., no evidence was found for an association between endometriosis and the risk of hypertension or preeclampsia in pregnancy.⁵¹ In our results, while the GHT rate was found to be significant in the endometriosis group compared to the non-endometriosis group, no difference was observed between the groups in terms of preeclampsia risk. The increased incidence of PROM in pregnancy in women with endometriosis is associated with a physical/microbial inflammatory process that weakens the

fetal membranes and increases prostaglandins (PGs), which in turn lead to collagen degradation within the fetal membranes through the action of metalloproteinases (MMP-9) and collagenase.⁵² Increased levels of PG and inflammatory cytokines in women with endometriosis locally activate MMP-9 and matrix-degrading enzymes and are responsible for the invasiveness of lesions.^{31,53} In the study by Conti et al., the risk of PROM was significantly higher in primiparous women with endometriosis compared with the control group.⁵⁴ In the study by Harada et al., PROM was found to be significantly higher in Endometriosis group compared to non-endometriosis group who conceived naturally or received infertility treatment other than ART treatment.³² Although our study revealed that the rate of PROM was higher in the endometriosis group, it did not detect a significant result. Pérez-López et al.'s meta-analysis stated that endometriosis had no significant effect on the risk of GDM.⁵⁵ The study by Salmeri et al. revealed an increased risk of GDM in endometriosis and with a possible progressive effect in more advanced stages of the disease.⁵⁶ In addition to the different results presented in the literature, our study did not detect a clear relationship between endometriosis and GDM in the data we obtained. This difference in the studies may be related to the way endometriosis is diagnosed and the degree to which the severity of endometriosis is different. In Lalani et al.'s meta-analysis, women with endometriosis were more likely to be admitted to the NICU.³⁷ Similarly, in the meta-analysis by Horton et al., it was reported that the probability of being admitted to the NICU was higher for women with Endometriosis.⁵⁷ Our current data are consistent with the literature results, and it has been concluded that NICU demand is significantly higher in the presence of endometriosis. In the evaluation of Breintoft et al., the cesarean ratio was found to be higher in the endometriosis group than in the control group.³⁸ Similarly, in present study, the cesarean ratio was found to be higher in the endometriosis group. The reason for this may be the decision to perform cesarean section due to the secondary outcome of perinatal complications. However, the general increase in our cesarean rates compared to the litera-

ture has become a public health issue that needs to be examined in depth. We consider our study as one of the rare multicentric cohort studies in our country that evaluates obstetric and neonatal outcomes for women with endometriosis on a large scale and multifactorially. Our study had a consistent methodology and considered the way patients conceived, and we believe this may have an independent effect on the outcomes of interest. In endometriosis, surgery and histology continue to be the gold standard diagnostic techniques internationally. Most of the patients in the endometriosis group were diagnosed with endometriosis before pregnancy, which can be considered as an advantage of this study. We accept, specifically in our study, that women in the endometriosis group diagnosed only by ultrasound did not receive surgical confirmation of endometriosis and that there may be cases of endometriosis that were misdiagnosed on ultrasound. The fact that endometriosis was not surgically confirmed in all patients in our study may be considered a limitation. However, laparoscopy is no longer accepted as a diagnostic reference standard for endometriosis and is now recommended only in women with persistent symptoms and negative imaging results or in those who have failed empirical treatment.⁵⁸ We acknowledge that we may have missed detecting endometriosis in some women in the group without endometriosis, particularly in women with peritoneal disease. Peritoneal endometriosis is common, not always detected on pelvic ultrasound, and may be found incidentally at laparoscopy.⁵⁹ It is clear that including only women with surgical diagnosis would be a more robust method for screening and defining disease subtypes. However, women with endometriosis are increasingly being treated conservatively, and including only those with surgical diagnosis would have limited the population studied to women with symptomatic disease or those who elected surgery. There is a possibility that there were also women with mild and minimal endometriosis in the group of patients without endometriosis, and when evaluating the findings of our study, careful interpretation is necessary regarding women with mild and minimal disease.

CONCLUSION

Perinatal and neonatal outcomes resulting from endometriosis depend on multifactorial factors. We believe that the sample size in our study population may lead to associations that are not statistically significant and therefore prospective and large population-based studies or meta-analyses are needed to provide meaningful results and clarify possible risks.

Source of Finance

During this study, no financial or spiritual support was received neither from any pharmaceutical company that has a direct connection with the research subject, nor from a company that pro-

vides or produces medical instruments and materials which may negatively affect the evaluation process of this study.

Conflict of Interest

No conflicts of interest between the authors and / or family members of the scientific and medical committee members or members of the potential conflicts of interest, counseling, expertise, working conditions, share holding and similar situations in any firm.

Authorship Contributions

Idea/Concept: Ufuk Atlıhan, Selçuk Erkalıç; **Design:** Onur Yavuz; **Control/Supervision:** Ferruh Acet; **Data Collection and/or Processing:** Can Ata; **Analysis and/or Interpretation:** Teyfik Berk Bildacı; **Literature Review:** Ufuk Atlıhan; **Writing the Article:** Ufuk Atlıhan, Burak Ersak; **Critical Review:** Hüseyin Aytuğ Aşar.

REFERENCES

1. Atlıhan U, Ertan B, Özgözen E, Güney M. Comparison of Patients with Adenomyosis Detected in Hysterectomy Material and Patients with Other Benign Pathologies: Retrospective Study. *Turkish Journal of Reproductive Medicine and Surgery*. 2024;8(2):54-62.
2. Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, Hummelshoj L, Prentice A, Saridogan E. ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod*. 2005;20:2698-704.
3. Atlıhan U, Yavuz O, Aşar HA, Ata C, Erkalıç S, Bildacı TB. Vitamin D evaluation in adenomyosis: A retrospective cross-sectional study. *Turk J Obstet Gynecol*. 2024;21(2):98-103.
4. Buck Louis GM, Hediger ML, Peterson CM, Croughan M, Sundaram R, Stanford J, et al. ENDO Study Working Group. Incidence of endometriosis by study population and diagnostic method: the ENDO study. *Fertil Steril*. 2011;96(2):360-5.
5. Bean E, Naftalin J, Horne A, Saridogan E, Cutner A, Jurkovic D. Prevalence of deep and ovarian endometriosis in early pregnancy: ultrasound diagnostic study. *Ultrasound Obstet Gynecol*. 2022;59(1):107-113.
6. Naftalin J, Hoo W, Pateman K, Mavrelis D, Holland T, Jurkovic D. How common is adenomyosis? A prospective study of prevalence using transvaginal ultrasound in a gynaecology clinic. *Hum Reprod*. 2012;27(12):3432-9.
7. May KE, Villar J, Kirtley S, Kennedy SH, Becker CM. Endometrial alterations in endometriosis: a systematic review of putative biomarkers. *Hum Reprod Update*. 2011;17(5):637-53.
8. Brosens I, Pijnenborg R, Benagiano G. Defective myometrial spiral artery remodelling as a cause of major obstetrical syndromes in endometriosis and adenomyosis. *Placenta*. 2013;34(2):100-5.
9. Brosens I, Brosens JJ, Fusi L, Al-Sabbagh M, Kuroda K, Benagiano G. Risks of adverse pregnancy outcome in endometriosis. *Fertil Steril*. 2012;98(1):30-5.
10. Fernando S, Breheny S, Jaques AM, Halliday JL, Baker G, Healy D. Preterm birth, ovarian endometriomas, and assisted reproduction technologies. *Fertil Steril*. 2009;91(2):325-30.
11. Juang CM, Chou P, Yen MS, Twu NF, Horng HC, Hsu WL. Adenomyosis and risk of preterm delivery. *BJOG*. 2007;114(2):165-9.
12. Stephansson O, Kieler H, Granath F, Falconer H. Endometriosis, assisted reproduction technology, and risk of adverse pregnancy outcome. *Hum Reprod*. 2009;24(9):2341-7.
13. Omland AK, Abyholm T, Fedorcsák P, Ertzeid G, Oldereid NB, Bjercke S, et al. Pregnancy outcome after IVF and ICSI in unexplained, endometriosis-associated and tubal factor infertility. *Hum Reprod*. 2005;20(3):722-7.
14. Vercellini P, Parazzini F, Pietropaolo G, et al. Pregnancy outcome in women with peritoneal, ovarian and rectovaginal endometriosis: a retrospective cohort study. *BJOG* 2012;119:1538-43.
15. Benaglia L, Bermejo A, Somigliana E, Scardueli C, Ragni G, Fedele L, Garcia-Velasco JA. Pregnancy outcome in women with endometriomas achieving pregnancy through IVF. *Hum Reprod*. 2012;27(6):1663-7.
16. Mekaru K, Masamoto H, Sugiyama H, Asato K, Heshiki C, Kinjyo T, Aoki Y. Endometriosis and pregnancy outcome: are pregnancies complicated by endometriosis a high-risk group? *Eur J Obstet Gynecol Reprod Biol*. 2014;172:36-9.
17. Shmueli A, Salman L, Hirsch L, Ashwal E, Hadar E, Wiznitzer A, et al. Obstetrical and neonatal outcomes of pregnancies complicated by endometriosis. *J Matern Fetal Neonatal Med*. 2019;32(5):845-50.
18. Saraswat L, Ayansina DT, Cooper KG, Bhattacharya S, Milligkos D, Horne AW, Bhattacharya S. Pregnancy outcomes in women with endometriosis: a national record linkage study. *BJOG*. 2017;124(3):444-52.
19. Lin H, Leng JH, Liu JT, Lang JH. Obstetric outcomes in Chinese women with endometriosis: a retrospective cohort study. *Chin Med J (Engl)*. 2015;128(4):455-8.
20. Kuivasaari-Pirinen P, Raatikainen K, Hippeläinen M, Heinonen S. Adverse Outcomes of IVF/ICSI Pregnancies Vary Depending on Aetiology of Infertility. *ISRN Obstet Gynecol*. 2012;2012:451915.
21. Harmsen MJ, Van den Bosch T, de Leeuw RA, Dueholm M, Exacoustos C, Valentin L, et al. Consensus on revised definitions of Morphological Uterus Sonographic Assessment (MUSA) features of adenomyosis: results of modified Delphi procedure. *Ultrasound Obstet Gynecol*. 2022;60(1):118-31.
22. Atlıhan U. Myoma uteriye yaklaşım. Aslan M, Sarıtaş AG, Aksoy S, Atlıhan U, editörler. *Sağlıkta Güncel Gelişmeler Işığında Teorik ve Klinik Yaklaşımlar-4*. Bursa: Ekin Yayınevi; 2023. p.50-71.

23. Pfeifer SM, Attaran M, Goldstein J, Lindheim SR, Petrozza JC, Rackow BW, et al. ASRM müllerian anomalies classification 2021. *Fertil Steril*. 2021;116(5):1238-52.
24. Guerriero S, Condous G, van den Bosch T, Valentin L, Leone FP, Van Schoubroeck D, et al. Systematic approach to sonographic evaluation of the pelvis in women with suspected endometriosis, including terms, definitions and measurements: a consensus opinion from the International Deep Endometriosis Analysis (IDEA) group. *Ultrasound Obstet Gynecol*. 2016;48(3):318-32.
25. American Diabetes Association. 2. Classification and Diagnosis of Diabetes: Standards of Medical Care in Diabetes-2020. *Diabetes Care*. 2020;43(Suppl 1):S14-S31.
26. ACOG Practice Bulletin No. 202: Gestational Hypertension and Preeclampsia. *Obstet Gynecol*. 2019;133(1):1.
27. Gordijn SJ, Beune IM, Thilaganathan B, Papageorghiou A, Baschat AA, Baker PN, et al. Consensus definition of fetal growth restriction: a Delphi procedure. *Ultrasound Obstet Gynecol*. 2016;48(3):333-9.
28. Stafford I, Diddy GA, Clark SL, Belfort MA. Visually estimated and calculated blood loss in vaginal and cesarean delivery. *Am J Obstet Gynecol*. 2008;199(5):519.e1-7.
29. Brosens I, Pijnenborg R, Vercruyse L, Romero R. The "Great Obstetrical Syndromes" are associated with disorders of deep placentation. *Am J Obstet Gynecol*. 2011;204(3):193-201.
30. Pizzo A, Salmeri FM, Ardita FV, Sofo V, Tripepi M, Marsico S. Behaviour of cytokine levels in serum and peritoneal fluid of women with endometriosis. *Gynecol Obstet Invest*. 2002;54(2):82-7.
31. Rakhila H, Carli C, Daris M, Lemyre M, Leboeuf M, Akoum A. Identification of multiple and distinct defects in prostaglandin biosynthetic pathways in eutopic and ectopic endometrium of women with endometriosis. *Fertil Steril*. 2013;100(6):1650-9.e1-2.
32. Harada T, Taniguchi F, Onishi K, Kurozawa Y, Hayashi K, Harada T; Japan Environment & Children's Study Group. Obstetrical Complications in Women with Endometriosis: A Cohort Study in Japan. *PLoS One*. 2016;11(12):e0168476.
33. Farland LV, Davidson S, Sasamoto N, Horne AW, Missmer SA. Adverse Pregnancy Outcomes in Endometriosis - Myths and Realities. *Curr Obstet Gynecol Rep*. 2020;9(1):27-35.
34. Petraglia F, Arcuri F, de Ziegler D, Chapron C. Inflammation: a link between endometriosis and preterm birth. *Fertil Steril*. 2012;98(1):36-40.
35. Kunz G, Beil D, Huppert P, Leyendecker G. Structural abnormalities of the uterine wall in women with endometriosis and infertility visualized by vaginal sonography and magnetic resonance imaging. *Hum Reprod*. 2000;15(1):76-82.
36. Aguilar HN, Mitchell BF. Physiological pathways and molecular mechanisms regulating uterine contractility. *Hum Reprod Update*. 2010;16(6):725-44.
37. Lalani S, Choudhry AJ, Firth B, Bacal V, Walker M, Wen SW, et al. Endometriosis and adverse maternal, fetal and neonatal outcomes, a systematic review and meta-analysis. *Hum Reprod*. 2018;33(10):1854-65.
38. Breintoft K, Pinnerup R, Henriksen TB, Rytter D, Ulbjerg N, Forman A, Arendt LH. Endometriosis and Risk of Adverse Pregnancy Outcome: A Systematic Review and Meta-Analysis. *J Clin Med*. 2021;10(4):667.
39. Exacoustos C, Lauriola I, Lazzari L, De Felice G, Zupi E. Complications during pregnancy and delivery in women with untreated rectovaginal deep infiltrating endometriosis. *Fertil Steril*. 2016;106(5):1129-35.e1.
40. Farella M, Chanavaz-Lacheray I, Verspick E, Merlot B, Klapczynski C, Henttier C, et al. Pregnancy outcomes in women with history of surgery for endometriosis. *Fertil Steril*. 2020;113(5):996-1004.
41. Aris A. A 12-year cohort study on adverse pregnancy outcomes in Eastern Townships of Canada: impact of endometriosis. *Gynecol Endocrinol*. 2014;30(1):34-7.
42. Mitri F, Casper RF. Endometriosis and mild bleeding disorders. *Fertil Steril*. 2015;103(4):886-7.e1.
43. Katorza E, Soriano D, Stockheim D, Mashiach R, Zolti M, Seidman DS, et al. Severe intraabdominal bleeding caused by endometriotic lesions during the third trimester of pregnancy. *Am J Obstet Gynecol*. 2007;197(5):501.e1-4.
44. Barbieri M, Somigliana E, Oneda S, Ossola MW, Acaia B, Fedele L. Decidualized ovarian endometriosis in pregnancy: a challenging diagnostic entity. *Hum Reprod*. 2009;24(8):1818-24.
45. Canlorbe G, Goubin-Versini I, Azria E, Abdul-Razak R, Muray J-M. La deciduose ectopique: variabilites de presentation pendant la grossesse et diagnostics differentiels. *Gynecol Obstet Fertil*. 2012;40:235-40.
46. Bean E, Cutner A, Saridogan E, Wong M, Naftalin J, Jurkovic D. Hemoperitoneum as a precursor of deep pelvic endometriosis: prospective cohort study. *Ultrasound Obstet Gynecol*. 2019;54(3):389-94.
47. Lafleur N, Wei SQ, Bilodeau-Bertrand M, Auger N. Association of Endometriosis and Severe Maternal Morbidity. *Obstet Gynecol*. 2022;140(6):1008-16.
48. Glavind MT, Forman A, Arendt LH, Nielsen K, Henriksen TB. Endometriosis and pregnancy complications: a Danish cohort study. *Fertil Steril*. 2017;107(1):160-6.
49. Chen I, Lalani S, Xie RH, Shen M, Singh SS, Wen SW. Association between surgically diagnosed endometriosis and adverse pregnancy outcomes. *Fertil Steril*. 2018;109(1):142-7.
50. Farland LV, Prescott J, Sasamoto N, Tobias DK, Gaskins AJ, Stuart JJ, et al. Endometriosis and Risk of Adverse Pregnancy Outcomes. *Obstet Gynecol*. 2019;134(3):527-36.
51. Hadfield RM, Lain SJ, Raynes-Greenow CH, Morris JM, Roberts CL. Is there an association between endometriosis and the risk of pre-eclampsia? A population based study. *Hum Reprod*. 2009;24(9):2348-52.
52. Vadillo-Ortega F, Estrada-Gutiérrez G. Role of matrix metalloproteinases in preterm labour. *BJOG*. 2005;112 Suppl 1:19-22.
53. Pitsos M, Kanakas N. The role of matrix metalloproteinases in the pathogenesis of endometriosis. *Reprod Sci*. 2009;16(8):717-26.
54. Conti N, Cevenini G, Vannuccini S, Orlandini C, Valensise H, Gervasi MT, et al. Women with endometriosis at first pregnancy have an increased risk of adverse obstetric outcome. *J Matern Fetal Neonatal Med*. 2015;28(15):1795-8.
55. Pérez-López FR, Martínez-Domínguez SJ, Viñas A, Pérez-Tambo R, Lafita A, Lajusticia H, et al. Health Outcomes and Systematic Analyses (HOUSSAY) Project. Endometriosis and gestational diabetes mellitus risk: a systematic review and meta-analysis. *Gynecol Endocrinol*. 2018;34(5):363-9.
56. Salmeri N, Li Piani L, Cavoretto PI, Somigliana E, Viganò P, Candiani M. Endometriosis increases the risk of gestational diabetes: a meta-analysis stratified by mode of conception, disease localization and severity. *Sci Rep*. 2023;13(1):8099.
57. Horton J, Sterrenburg M, Lane S, Maheshwari A, Li TC, Cheong Y. Reproductive, obstetric, and perinatal outcomes of women with adenomyosis and endometriosis: a systematic review and meta-analysis. *Hum Reprod Update*. 2019;25(5):592-632.
58. Becker CM, Bokor A, Heikinheimo O, Horne A, Jansen F, Kiesel L, et al. ESHRE Endometriosis Guideline Group. ESHRE guideline: endometriosis. *Hum Reprod Open*. 2022;2022(2):hoac009.
59. Hager M, Wenzl R, Riesenhuber S, Marschalek J, Kuessel L, Mayrhofer D, et al. The Prevalence of Incidental Endometriosis in Women Undergoing Laparoscopic Ovarian Drilling for Clomiphene-Resistant Polycystic Ovary Syndrome: A Retrospective Cohort Study and Meta-Analysis. *J Clin Med*. 2019;8(8):1210.